






























Soils Legend

	1016 - Udorthents, loamy
	102B - Clarion loam, 2 to 6 percent slopes
	106B - Lester loam, 2 to 6 percent slopes
	106C2 - Lester loam, 6 to 12 percent slopes, eroded
	1081 - Minneiska-Abscota complex, occasionally flooded
	109 - Cordova clay loam
	112 - Harps clay loam
	118 - Crippin loam
	1833 - Coland clay loam, occasionally flooded
	1834 - Coland clay loam, frequently flooded
	1901B - Lester-Le Sueur complex, 1 to 6 percent slopes
	239 - Le Sueur clay loam
	27B - Dickinson loam, 2 to 6 percent slopes
	336 - Delft clay loam
	386 - Okoboji mucky silty clay loam
	86 - Canisteo clay loam
	920B - Clarion-Hawick complex, 3 to 6 percent slopes
	921C2 - Clarion-Storden complex, 6 to 12 percent slopes, eroded
	944B - Lester-Hawick complex, 2 to 6 percent slopes
	945D2 - Lester-Storden complex, 12 to 18 percent slopes, eroded
	946 - Nicollet-Linder complex
	978 - Cordova-Rolfe complex
	L107A - Canisteo-Glencoe, depressional complex, 0 to 2 percent slopes
	L13A - Klossner muck, depressional, 0 to 1 percent slopes
	L163A - Okoboji silty clay loam, depressional, 0 to 1 percent slopes
	L83A - Webster clay loam, 0 to 2 percent slopes
	L84A - Glencoe clay loam, depressional, 0 to 1 percent slopes
	L85A - Nicollet clay loam, 1 to 3 percent slopes
	W -